ΑD	١		

Award Number: W81XWH-14-1-0011

TITLE:

Identification of Prostate Cancer-Specific microDNAs

PRINCIPAL INVESTIGATOR:

Yin-Yuan Mo

CONTRACTING ORGANIZATION:

University of Mississippi Medical Center Jackson, MS 39216

REPORT DATE: December 2014

TYPE OF REPORT:

Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

Form Approved REPORT DOCUMENTATION PAGE OMB No. 0704-0188 Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. 1. REPORT DATE 2. REPORT TYPE 3. DATES COVERED December 2014 Annual 25 Nov 2013 - 24 Nov 2014 4. TITLE AND SUBTITLE 5a. CONTRACT NUMBER Identification of Prostate Cancer-Specific microDNAs 5b. GRANT NUMBER W81XWH-14-1-0011 **5c. PROGRAM ELEMENT NUMBER** 6. AUTHOR(S) 5d. PROJECT NUMBER Yin-Yuan Mo 5e. TASK NUMBER 5f. WORK UNIT NUMBER E-Mail: ymo@umc.edu 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 8. PERFORMING ORGANIZATION REPORT NUMBER University of Mississippi Medical Center 2500 State Street, G652 Jackson, MS 39216 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSOR/MONITOR'S ACRONYM(S) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012 11. SPONSOR/MONITOR'S REPORT NUMBER(S) 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited 13. SUPPLEMENTARY NOTES 14. ABSTRACT Emerging evidence has suggested that eukaryotic cells can express a special group of extrachromosomal circular DNAs (eccDNAs), called microDNAs. Unlike previously reported eccDNAs, microDNAs are relatively small in length, map to unique DNA sequence, and arise from genes, mostly likely resulting from microdeletions. Since they are usually in a circular form, they are more resistant to exonuclease than linear DNAs and can be stably present in the cells or even possibly in the circulating system. Therefore, overall goal of this application is to determine whether prostate cancer cells express such microDNAs which can be used to serve as valuable biomarkers for prostate cancer diagnosis or prognosis. 15. SUBJECT TERMS microRNA, lncRNA, biomarker

17. LIMITATION

OF ABSTRACT

UU

18. NUMBER

OF PAGES

7

19a. NAME OF RESPONSIBLE PERSON

19b. TELEPHONE NUMBER (include area

USAMRMC

16. SECURITY CLASSIFICATION OF:

b. ABSTRACT

c. THIS PAGE

a. REPORT

Table of Contents

	<u>Page</u>
Introduction	1
Body	1
Key Research Accomplishments	4
Reportable Outcomes	4
Conclusion	4
References	N/a
Appendices	N/a

Introduction

MicroDNAs are a special group of extrachromosomal circular DNAs (eccDNAs) derived from chromosomal repetitive sequences, intermediates of mobile elements or viral genomes. Unlike most eccRNAs, microDNAs are small, map to unique DNA sequence, and arise from genes, mostly likely resulting from microdeletions. Since they are usually in a circular form, they are resistant to exonuclease and more stable present in the cells or even possibly in the circulating system. We hypothesize that prostate cancer may exploit this mechanism for its own advantage and thus may express a very different microDNA pattern from normal prostate tissue. This different pattern can be detected by currently advanced technology such as deep sequencing. Therefore, overall goal of this application is to determine whether prostate cancer cells express specific microDNAs which may serve a potential biomarkers for prostate cancer diagnosis or prognosis.

Body




Fig. 1Amplification of DNA by multiple displacement amplification (MDA).

Little is known about
microDNAs, and it is not
clear whether prostate
cancer cells carry
potential microDNAs.
Thus our goal is to
demonstrate the
existence of microDNAs
in prostate cancer. We

adopted multiple displacement amplification (MDA) with random primers for enriched circular DNA by rolling circle amplification (RCA) (Fig. 1) and then amplified DNA fragments were subject to deep sequencing.

Sequence	NO of Reads
seq 1	184
seq 2	133
seq 3	2407
seq 4	166
seq 5	126
seq 6	221
seq 7	1388
seq 8	108
seq 9	139
seq 10	158
seq 11	968
seq 12	115
seq 13	460
seq 14	391
seq 15	364
seq 16	157
seq 17	634
seq 18	438
seq 19	368
seq 20	529
seq 21	209
seq 22	171
seq 23	271
seq 24	110
seq 25	254
seq 26	187
seq 27	109
seq 28	191
seq 29	103
seq 30	188
seq 31	125





Fig.2 Identification of potential microDNAs from prostate cancer.



Deep sequencing of the amplified DNA fragments identified several potential microDNA sequences. Among them, 31 sequences were listed in Fig. 2. The detailed sequences of 4 clones were shown in Fig. 3. All 4 clones were within 1kb in length. Of interest, clone 7 carries AA, AT or TT dinucleotides, a feature of microDNAs; In addition, its GC content is relatively high (58.4%), another feature of microDNAs.

>LNCAP2-7 (286bp)

GTATAATGTGGTGGCCAGGTGCAGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCG AGGTGGGCGGATCACGAGGTCAGGAGATCGAGACCATCCTGGCTAACATGGTGAAACCCC GTCTCTACTAAAAATACAAAAAATTAGCCGGGCGTGGTGGTGGGCACCTGTAGTCCCAGCTA CTCGGGAGGCTGAGGCAGGAGAATGGCGTGAACCCGGGAGGCGGAGCTTGCAGTGAGC CGAGATCACGCCACTGCACTCCAGCCTGGGTGACAGAGTGAGACA

Fig. 4 This clone carries multiple AA, AT or TT dinucleotides, a feature of microDNAs, in addition to relatively high GC content.

Identification of such potential microDNAs is critical to the next stage of investigation, i.e., to determine whether such potential microDNAs are present in blood/serum of prostate cancer patients such that they may serve as biomarkers for prostate cancer. Future work will determine whether the potential microDNAs will impact prostate tumor cell growth and invasion. We will also determine whether they are differently present in prostate cancer patients.

Key Research Accomplishments

- We identified several potential microDNAs from prostate cancer cells through multiple displacement amplification.
- Clone #7 is the top candidate which has been cloned in an expression vector and it
 will be tested in prostate cancer cell lines for its effect on tumor cell growth and
 invasion
- We will determine whether microDNAs such as clone #7 sequence are present in blood/serum samples of prostate cancer patients and test their potential as biomarkers for prostate cancer.

Reportable Outcomes

Not yet.

Conclusions

Deep sequencing of MDA samples from prostate cancer cells has identified 31 potential microDNA candidates. Clone #7 is a top candidate based on size, dinucleotide repeats and high GC content. Furthermore, we have cloned this sequence. We are currently determining whether ectopic expression of clone #7 will impact prostate tumor cell growth and invasion.